

# Additional Experimental Proof of the Existence of the Second Flavor of Hydrogen Atoms from the Analysis of Experimental Cross-Sections of Charge Exchange between Hydrogen Atoms and Protons

E. OKS

*Physics Department, 380 Duncan Drive, Auburn University, Auburn, AL 36849, USA*

**ABSTRACT:** There is a noticeable discrepancy between the experimental and previous theoretical results on cross-sections of charge exchange between hydrogen atoms and low energy protons. The experimental cross-sections were systematically slightly higher – beyond the error margins – than the theoretical predictions. In the present paper we analyze whether this discrepancy can be eliminated or at least reduced by using in calculations the Second Flavor of Hydrogen Atoms (SFHA). We demonstrate that for the SFHA the corresponding cross-section is noticeably larger than for the usual hydrogen atoms. We show that the allowance for the SFHA does bring the theoretical cross-sections in a noticeably better agreement with the corresponding experiments within the experimental error margins. This seems to constitute yet another proof from atomic experiments that the SFHA is present within the mixture of hydrogen atoms. In combination with the first corresponding proof from the analysis of atomic experiments (concerning the distribution of the linear momentum in the ground state of hydrogen atoms), as well as with the astrophysical evidence from two different kinds of observations (the anomalous absorption of the redshifted 21 cm radio line from the early Universe and the smoother distribution of dark matter than predicted by the standard cosmology), the results of the present paper reinforce the status of the SFHA as the candidate for dark matter or at least for a part of it.

**Keywords:** charge exchange; second flavor of hydrogen atoms; dark matter; Stark effect

## 1. INTRODUCTION

There is a noticeable discrepancy between the experimental and previous theoretical results on cross-sections of charge exchange between hydrogen atoms and low energy protons (down to the energy  $\sim 10$  eV). Experiments by Fite et al [1], Fite et al [2], and Belyaev et al [3] revealed that the experimental cross-sections were systematically slightly higher – beyond the error margins – than the theoretical predictions. The source of this discrepancy was stated as unknown [3]. Up to now there was no attention paid to his discrepancy, to the best of our knowledge.

In paper [4] we showed analytically that there is a slight difference in cross-sections of charge exchange between the usual hydrogen atoms and protons in comparison to charge exchange between the Second Flavor of Hydrogen Atoms (SFHA) and protons. The SFHA was discovered theoretically, as well as by the analysis of atomic experiments, in paper [5].

The basis of the theoretical discovery in paper [5] was a fresh analysis of the Dirac equation for hydrogen atoms. There are two solutions of the Dirac equation at relatively small distance from the proton: the regular solution and the singular solution, the latter being usually rejected. In paper [5] it was shown that at the proton boundary, the singular solution outside the proton can be tailored with the regular solution inside the proton for the ground state, so that the singular solution outside the proton is legitimate for the ground state. Using this fact, the author of paper [5] eliminated a huge discrepancy – by many orders of magnitude – between the experimental and theoretical distributions of the linear momentum in the ground state of hydrogen atoms. This constituted the first experimental evidence of the existence of the SFHA.

In the later paper [6] it was shown that the singular solution of the Dirac equation outside the proton is legitimate not only for the ground state, but for all discrete and continuous states of the zero angular momentum: for all S-states. This kind of hydrogen atoms, having only S-states and described by the singular solution of the Dirac equation outside the proton, was called the “second flavor” in paper [7] – by the analogy with the flavors of quarks.

By now there are also two kinds of the astrophysical evidence of the existence of the SFHA. First, the SFHA eliminated a factor of two discrepancy between the absorption signal at the redshifted 21 cm radio line, recently observed by Bowman et al [8] and the predictions of the standard cosmology – this was shown in paper [6]. Second, the SFHA explained the recent observations by Jeffrey et al [9], where they found that the distribution of dark matter in the Universe is noticeably smoother than predictions based on Einstein’s relativity – the explanation was provided in paper [10].

The most striking feature of the SFHA is that since they have only S-states, then due to the well-known selection rules, the SFHA does not interact with the electromagnetic radiation (except for the radiative transition at the 21 cm wavelength between the two hyperfine structure substates of the ground state). Thus, the SFHA remains “dark”. In combination with the above two kinds of the astrophysical evidence, this makes the SFHA a good candidate for dark matter or at least for a part of it.

The purpose of the present paper is to find out whether by using the SFHA, one can eliminate or at least reduce the aforementioned discrepancy between the experimental and theoretical cross-sections of charge exchange involving hydrogen atoms and low energy protons. For this purpose we extend the theory from paper [4] (where it was developed for excited states) specifically for the ground state – since the corresponding experiments dealt with hydrogen atoms in the ground state. We demonstrate that for the SFHA the corresponding cross-section is noticeably larger than for the usual hydrogen atoms. We show that the allowance for the SFHA does bring the theoretical cross-sections in the better agreement with the corresponding experiments within the experimental error margins.

## 2. CALCULATIONS AND THE COMPARISON WITH EXPERIMENTS

According to paper [11], classically the cross-section  $\sigma$  for the resonant transition of the electron from being associated with one ion to being associated with another ion is

$$\sigma = (8\pi/I^2)(1 - 0.8z^{2/5}), \quad z = v/(2I)^{1/2}. \quad (1)$$

Here  $v$  is the relative velocity of the colliding nuclei and  $I$  is the ionization potential from the particular atomic state. In the present paper we use atomic units, unless noted otherwise. Formula (1) is valid for relatively small velocities:

$$v \ll v_{\max} = (2I)^{1/2}. \quad (2)$$

We study the resonant charge exchange between a hydrogen atom in the ground state and an incoming proton, so that from now on,  $I$  is the ionization potential of hydrogen atoms from the ground state. For calculating the quantity  $I$ , we take into account the Stark shift (if any) of the ground state due to the electric field the proton separated by the distance  $R$  from the atom.

It should be emphasized upfront that only the energy levels of the usual hydrogen atoms experience the Stark shift. In distinction, the energy levels of the SFHA has *no Stark shift* in the field of the proton *in any order of the multipole expansion*, as explained in paper [4]. This fact leads to the difference in the corresponding cross-sections of charge exchange for collisions of protons with the usual hydrogen atom compared to the collision of protons with the SFHA, as calculated below.

For relatively large  $R$ , the ground state energy of *usual* hydrogen atoms can be represented in the form (according to Eq. (4.59) from book [12])

$$E_{\text{large}} = -1/2 - 9/(4R^4) - 15/(2R^6), \quad (3)$$

where we omitted higher order terms in this expansion; the subscript “large” signifies that this expansion is valid for relatively large  $R$ . So, the ionization potential for the ground state has the form:

$$I_{\text{large}} = 1/2 + 9/(4R^4) + 15/(2R^6). \quad (4)$$

In paper [11] it was noted that for the no-barrier-transition of the electron from one nucleus to another nucleus, at the midpoint between the two nuclei the interaction potential for the electron in the field of two nuclei should surpass the corresponding ionization potential. This condition translates in the following relation between the charge-exchange-effective distance  $R_0$  of the proton from the atom and the ionization potential [11]:

$$I = 4/R_0. \quad (5)$$

On substituting Eq. (5) in Eq. (1), we obtain:

$$\sigma = (\pi R_0^2/2)[1 - 0.8v^{2/5}(R_0/8)^{1/5}]. \quad (6)$$

Further, on substituting Eq. (5) in the left side of Eq. (4) (and omitting the subscript “0”) we get:

$$4/R = 1/2 + 9/(4R^4) + 15/(2R^6). \quad (7)$$

The relevant root of this equation is

$$R_{\text{large}} = 7.991. \quad (8)$$

(We remind that the subscript “large” here and below just refers to the fact that the results were obtained from the energy expansion for relatively large  $R$ .)

In distinction, for the SFHA for relatively large  $R$  the energy of the ground state is

$$E_{\text{large},2} = -1/2 \quad (9)$$

because the SFHA does not experience any Stark shift. (Here the number 2 in the subscript signifies the “second flavor”.) Therefore,

$$I_{\text{large}} = 1/2, \quad (10)$$

so that (according to Eq. (5))

$$R_{\text{large},2} = 8. \quad (11)$$

Now let us consider the corresponding situation for relatively small  $R$ . In this case, the ground state energy of *usual* hydrogen atoms can be represented in the form (according to Eq. (5.13) from book [12])

$$E_{\text{small}} = -2 + 8R^2/3 - 16R^3/3, \quad (12)$$

where we omitted higher order terms in this expansion; the subscript “small” signifies that this expansion is valid for relatively small  $R$ . So, the ionization potential for the ground state has the form:

$$I_{\text{small}} = 2 - 8R^2/3 + 16R^3/3. \quad (13)$$

Further, on substituting Eq. (5) in the left side of Eq. (13) (and omitting the subscript “0”) we get:

$$4/R = 2 - 8R^2/3 + 16R^3/3. \quad (14)$$

The relevant root of this equation is

$$R_{\text{small}} = 0.953. \quad (15)$$

(We remind that the subscript “small” here and below just refers to the fact that the results were obtained from the energy expansion for relatively small  $R$ .)

In distinction, for the SFHA for relatively small  $R$  the energy of the ground state is

$$E_{\text{small},2} = -2 \quad (16)$$

because the energy levels of the SFHA do not shift in the electric field. (Here the number 2 in the subscript signifies

the “second flavor”.) Therefore,

$$I_{\text{small},2} = 2, \quad (17)$$

so that (according to Eq. (5))

$$R_{\text{small},2} = 2. \quad (18)$$

For calculating the cross-sections of the charge exchange by using Eq. (6), it is necessary to add the corresponding contributions from both channels, i.e., from both the “large R” case and the “small R” case. For obtaining the ratio of the resonant charge exchange cross-section from the ground state of the SFHA  $\sigma_{\text{SFHA}}$  to the corresponding result  $\sigma_{\text{usual}}$  for the usual hydrogen atoms in the simplest form – just to get the message across – we will consider the limit of  $v$  approaching zero. In this limit, the ratio  $\sigma_{\text{SFHA}}/\sigma_{\text{usual}}$  simplifies to:

$$\sigma_{\text{SFHA}}/\sigma_{\text{usual}} = (R_{\text{large},2}^2 + R_{\text{small},2}^2)/(R_{\text{large},2}^2 + R_{\text{small},2}^2). \quad (19)$$

On substituting in Eq. (19) the data from Eqs. (8), (11), (15), and (18), we finally obtain:

$$\sigma_{\text{SFHA}}/\sigma_{\text{usual}} = 1.05, \quad (20)$$

so that  $\sigma_{\text{SFHA}}$  is by 5% greater than  $\sigma_{\text{usual}}$ .

Let us find out whether this result sufficiently improves the comparison with the corresponding experiments. The most precise experiment on the resonant charge exchange between hydrogen atoms in the ground state and relatively low energy protons was performed by Fite et al [2]. Figure 1 presents:

- 1) the experimental cross-sections – circles, some with error margins – reproduced from Fig. 2 of paper [2];
- 2) the experimental cross-sections – crosses (with error margins) connected by the solid line – from the earlier measurements by Fite et al [1], reproduced from Fig. 2 of paper [2];
- 3) the theoretical cross-sections – filled circles connected by the solid line – calculated by Dalgarno and Yadav [13], reproduced from Fig. 2 of paper [2];
- 4) the theoretical cross-sections for the case of the SFHA – the dashed line – from the present calculations .

It is seen that the theoretical cross-sections for the case of the SFHA demonstrate a noticeably better agreement with the corresponding experimental cross-sections.




Fig. 1. Comparison of the experimental cross-sections of the resonant charge exchange between hydrogen atoms and low energy protons with the corresponding theoretical cross-sections (the cross-sections are in units of  $10^{-16}$  cm<sup>2</sup>). Circles (some with error margins) – experiment reported in paper [2]; crosses (with error margins) connected by the solid line – experiment reported in paper [1]; filled circles connected by the solid line – theory from paper [13], allowing only for the usual hydrogen atoms; the dashed line – theory from the present paper, allowing for the SFHA.

### 3. CONCLUSIONS

We analyzed whether the allowance for the SFHA can eliminate or at least reduce the noticeable discrepancy between the experimental and theoretical cross-sections of charge exchange involving hydrogen atoms and low energy protons: the discrepancy where the experimental cross-sections are systematically slightly higher than the corresponding theoretical cross-sections. We demonstrated that for the SFHA the theoretical cross-sections are noticeably greater than for the usual hydrogen atoms. We showed that the allowance for the SFHA leads to a noticeably better agreement with the experiments: the agreement with experiments within the experimental error margins.

This seems to constitute yet another proof from atomic experiments that the SFHA is present within the mixture of hydrogen atoms. In combination with the first corresponding proof from the analysis of atomic experiments (presented in paper [6]), as well as with the astrophysical evidence from two different kinds of observations [9, 11], the results of the present paper reinforce the status of the SFHA as the candidate for dark matter or at least for a part of it.

Compared to other explanations of dark matter effects, the SFHA is favored by the Occam razor principle. Indeed, it is based on the standard quantum mechanics (the Dirac equation), whereas other hypotheses either resort to mysterious, never discovered particles beyond the Standard Model or require significant changes of the existing physical laws.

### References

1. Fite, W.L.; Smith, A.C.H.; Stebbings, R.F. Charge Transfer in Collisions Involving Symmetric and Asymmetric Resonance. *Proceedings of the Royal Society*, **1962**, *A268*, 527-536.
2. Fite, W.L.; Brackmann, R.T.; Snow, W.R. Charge Exchange in Proton-Hydrogen-Atom Collisions. *Phys. Rev.* **1958**, *112*, 1161-1169.
3. Belyaev, V.A.; Brezhnev, B.G.; Erastov, E.M. Resonance Charge Exchange of Protons and Deuterons at Low Energies. *Soviet Physics JETP* **1967**, *25*, 777-782.
4. Oks, E. Classical Description of Resonant Charge Exchange Involving the Second Flavor of Hydrogen Atoms. *Atoms* **2021**, *9*, 41.
5. Oks, E. High-Energy Tail of the Linear Momentum Distribution in the Ground State of Hydrogen Atoms or Hydrogen-like Ions. *J. Phys. B: At. Mol. Opt. Phys.*, **2001**, *34*, 2235-2243.
6. Oks, E. Alternative Kind of Hydrogen Atoms as a Possible Explanation of the Latest Puzzling Observation of the 21 cm Radio Line from the Early Universe. *Research in Astronomy and Astrophysics* **2020**, *20(7)*, 109.
7. Oks, E. Two Flavors of Hydrogen Atoms: a Possible Explanation of Dark Matter. *Atoms* **2020**, *8*, 33.
8. Bowman, J.D.; Rogers, A.E.E.; Monsalve, R.A.; Mozdzen, T.J.; Mahesh, N. An Absorption Profile Centred at 78 Megahertz in the Sky-Averaged Spectrum. *Nature* **2018**, *555*, 67-70.
9. Jeffrey, N.; Gatti, M.; Chang, C; et al. Dark Energy Survey Year 3 results: Curved-sky weak lensing mass map reconstruction. *Monthly Notices of the Royal Astronomical Society* **2021**, *505(3)*, 4626-4645.
10. Oks, E. DES Map Shows a Smoother Distribution of Matter than Expected: A Possible Explanation. *Research in Astronomy and Astrophysics* **2021**, *21(10)*, 241-245.
11. Smirnov, B.M. The Classical Theory of Resonant Charge Exchange. *Sov. Phys. JETP* **1971**, *32*, 670-672.
12. Komarov, I.V.; Ponomarev, L.I.; Slavyanov, S. Yu. *Spheroidal and Coulomb Spheroidal Functions*; Nauka: Moscow, 1976, in Russian.
13. Dalgarno, A.; Yadav, H.N. Electron Capture II: Resonance Capture from Hydrogen Atoms by Slow Protons. *Proc. Phys. Soc. (London)* **1953**, *A66*, 173.